ATTENTION

This document is provided for historical purposes only.

Documents contained in the Washington Department of Fish and Wildlife Document & Publication Archive may contain dated and/or incorrect information. The WDFW Document & Publication Archive is provided as a service to those interested in the history of fish and wildlife management in Washington State.






Serving Washington's

wildlife and peoplenow and in the

future.



Doug Piler Bustrations courtsey of South Carolina Wildlife and Marine Resources Department and MWTF

SEX & AGE OF WILD TURKEYS

Determining the sex and age of wild turkeys is easy when you know what to look for.

Breast feathers

Differences in the tips of breast feathers are one method of determining sex in turkeys. Breast feathers are banded on the tips; bands are buff on hens and jet black on toms ("gobblers"). The bands are narrow on juveniles (less than a year old) and grow wider with age.

Beard

Older mature gobblers usually have beards which average seven to ten inches in length. Many juvenile males taken during the fall hunting season have beards which are shorter than two inches. Juvenile hens rarely possess beards, but a small portion of the adult females may be bearded. Beards on hens seldom exceed five inches in length and are sparse compared with the beard of a mature tom.

Head

Hens and juvenile males have small, dark heads covered with black hairs and miniature black feathers. The wattles, fleshy lobes that hang from the throat or chin, are dark and uncolored. A mature tom has a larger head with more wattles and less feathering. A tom's head is pink to pale red when the bird is relaxed, changing to hues of scarlet-red, with a white cap while strutting. When fright-ened during this displaying activity a tom's head turns bright blue. This may be a sign that the bird has detected the presence of the hunter.

Sours

Mature toms have prominent spurs on the backs of the lower legs, which become longer and more pointed with age. On an old bird, spurs can exceed an inch in length. Hens and juvenile toms have only rudimentary, inconspicuous spurs.

Outer primary or flight feathers
Primary feathers are the stiff

Primary feathers are the stiff, barred, black and white feathers on the wings. The two outer primary feathers on each wing are retained for a year. The tips of these feathers on juvenile birds appear sharp-pointed and worn, while the ends of outer primaries on adult birds are rounded and bars extend to the outer tips. On juveniles, the bars are absent from the outer inch or two of the feather tip.

Tail feathers

On a juvenile bird, the two center feathers are longer than the other tail feathers. Tail feathers on adults are equal in length.

Weight

The weight of a turkey depends on its sex, age, condition and subspecies. An average mature tom weighs between 17 and 21 lbs. in late fall; a mature hen, 10 to 13 lbs. Juvenile toms weigh 12 to 16 lbs. and young hens about 7 to 9 lbs.


Signs

The middle toe on a gobbler leaves a track at least four and one-half inches long, while the track of a hen or juvenile tom is shorter. Droppings of a gobbler are about the size of the little finger, with a slight crook at one end. Hen droppings are much smaller and shaped like a corkscrew.

The Washington Department of Whitelife will provide agent apparatus to to all potential and exhaining employable with materials in read, even other, some demanders, religion, egg, teached status, restand depth details. It is a substitution of the status of the status


The department reaching Federal Ald for Ball, and subdite projections. The department to exhipse in Tills VI of the Olivi Fights Act of 1984 on Bealish Side of the Federal Ballotton Added 1973, which prohibits department too on the basis advance, order, endound origin or benefitup. If you believe you have been districtionable qualent in any department program, and by, or family, or if you work before belowerialism, when Tills VI or Society, for family, or if you work before belowerialism, should Tilly VI or Society St., Westbergin, D.C. 2004, or Wisselington Department of Wildelin, 100 M. Capital Way, Chample, Will 2004.

SEX & AGE OF WILD TURKEYS




Age Determined By 3 Outer Wing Feethers






Doug Pifer illustrations courtesy of National Wild Turkey Federation



Breast Feathers



Doug Piter illustrations courtesy of South Carolina Wildlife and Marine Resources Department and NWTF